

The cross-correlation of the CMB polarisation and the 21 cm line fluctuations from cosmic reionisation

Hiroiyuki Tashiro¹, Nabila Aghanim¹, Mathieu Langer¹,
Marian Douspis¹, and Saleem Zaroubi²

¹ *Institut d'Astrophysique Spatiale (IAS), Bâtiment 121, F-91405, Orsay (France);
Université Paris-Sud XI and CNRS (UMR 8617)*

² *Kapteyn Astronomical Institute, University of Groningen, Postbus 800, NL-9700AV, Groningen, (The Netherlands)*

27 February 2008

ABSTRACT

The cosmic microwave background (CMB) polarisation and the 21 cm line fluctuations are powerful probes of cosmological reionisation. We study how the cross-correlation between the CMB polarisation (E -modes) and the 21 cm line fluctuations can be used to gain further understanding of the reionisation history, within the framework of inhomogeneous reionisation. Since the E -mode polarisation reflects the amplitude of the quadrupole component of the CMB temperature fluctuations, the angular power spectrum of the cross-correlation exhibits oscillations at all multipoles. The first peak of the power spectrum appears at the scale corresponding to the quadrupole at the redshift that is probed by the 21 cm line fluctuations. The peak reaches its maximum value in redshift when the average ionisation fraction of the universe is about half. On the other hand, on small scales, there is a damping that depends on the duration of reionisation. Thus, the cross-correlation between the CMB polarisation and the 21 cm line fluctuations has the potential to constrain accurately the epoch and the duration of reionisation.

Key words: cosmology: theory – large-scale structure of universe

1 INTRODUCTION

The history of the cosmological reionisation is one of the open problems of modern cosmology. Questions like what causes the reionisation and how it proceeds are intimately related to the evolution of matter density fluctuations and to the formation of the first structures (Barkana & Loeb 2001). The lack of observational data makes it quite difficult to answer these questions today. Up to now, the available probes of the reionisation epoch are just a few, e.g. the Gunn-Peterson test (Gunn & Peterson 1965) and the cosmic microwave background (CMB) radiation, in particular its polarisation (Zaldarriaga 1997). The observation of the Gunn-Peterson effect, which probes the amount of neutral hydrogen, suggests that full reionisation was accomplished by $z \sim 6$ (Fan et al. 2006). The CMB polarisation, in turn, measures the optical depth of the Thomson scattering from the last scattering surface to the present epoch. The WMAP 3 years data provide an optical depth $\tau \sim 0.09$ (Page et al. 2007) suggesting that reionisation began around $z = 10$.

In addition to those two probes, the observation of fluctuations of the 21 cm line background is expected to be one of the most promising methods to study reionisation (Madau et al. 1997; Tozzi et al. 2000; Ciardi & Madau 2003). The 21 cm line corresponds to the energy of the hyperfine spin flip transition of neutral hydrogen atoms. After decoupling of the baryon temperature from the CMB temperature around $z \sim 300$, neutral hydrogen atoms absorb or emit a 21 cm line, depending on their temperature. Since the 21 cm line is redshifted by the cosmological expansion, we can obtain redshift slices of the Universe by choosing the frequency of observation. It is thus possible to follow the evolution of the intergalactic medium before and during reionisation directly. The 21 cm line observations have recently gained a lot of attention (Loeb & Zaldarriaga 2004;

Furlanetto et al. 2006; Tashiro & Sugiyama 2006; Cooray 2006) and several projects are designed for measuring the fluctuating line background (e.g. MWA¹, LOFAR², SKA³).

In the near future, we will therefore have different direct and complementary probes of reionisation. We thus expect that cross-correlations between them will provide even more information than their respective auto-correlations. This is the case especially for the cross-correlation between the CMB temperature anisotropies and the 21 cm line fluctuations on large scales ($\ell \sim 100$ – Alvarez et al. 2006; Adshead & Furlanetto 2007) and on small scales ($\ell > 1000$ – Salvaterra et al. 2005; Cooray 2004; Slosar et al. 2007). Since the observations of the 21 cm lines are expected to be dominated by foregrounds and noise (Jelic et al. 2008), calculating the cross-correlation is indeed a good way to extract information on reionisation. In particular on large scales, the 21 cm line fluctuations are correlated with the CMB Doppler temperature anisotropies which are due to the motion of ionised baryons. Alvarez et al. (2006) showed that the maximum amplitude of the cross-correlation is reached at the redshift when the ionised fraction is one half, which potentially provides a way to constrain double reionisation models (e.g. Cen 2003).

In the present study, we examine the cross-correlation between the CMB E -mode polarisation (a more effective probe than the temperature anisotropies) and the fluctuations of the 21 cm line background on large scales. Measuring E -mode polarisation is one of the present challenging observations for CMB experiments. However, projects like PLANCK and CMB-pol will provide precise CMB polarisation maps in the near future. Similarly, cosmological observation of 21 cm line background is still belongs to the future, but LOFAR and MWA are being completed and will soon provide us with maps of neutral hydrogen at high redshifts. Hence, it is useful to investigate what information on reionisation we can extract from the cross-correlation of those maps. Indeed, we show here that the cross-correlations are a potentially powerful tool to constrain the history of cosmic reionisation. This article is organised as follows. In Sec. II, we provide the analytic formula of the cross-correlation between the CMB E -mode polarisation and the fluctuations of the 21 cm line background. In Sec. III, we show and discuss the results of the cross-correlation. Sec. VI is devoted to the conclusions. Throughout the paper, we use WMAP values for the cosmological parameters, i.e. $h = 0.73$ ($H_0 = h \times 100 \text{ km/s/Mpc}$), $T_0 = 2.725 \text{ K}$, $h^2 \Omega_b = 0.0223$ and $h^2 \Omega_m = 0.128$ (Spergel et al. 2007) for a flat cosmology. We also set the speed of light to $c = 1$.

2 CROSS-CORRELATION BETWEEN CMB E -MODE POLARISATION AND 21 CM LINES

2.1 The CMB E -mode polarisation

The CMB polarisation is best described by the Stokes parameters, Q and U . Since the Stokes parameters have spin 2, they can be decomposed onto the plane waves and spin-2 spherical harmonics ${}_{\pm 2}Y_\ell^m$ (Hu & White 1997),

$$(Q \pm iU)(\eta, \mathbf{x}, \hat{\mathbf{n}}) = \sum_{\ell m} (-i)^\ell \sqrt{\frac{4\pi}{2\ell+1}} \int \frac{d^3 k}{(2\pi)^3} \left(E_\ell^{(m)} \pm i B_\ell^{(m)} \right) \exp(i\mathbf{k} \cdot \mathbf{x}) {}_{\pm 2}Y_\ell^m(\hat{\mathbf{n}}). \quad (1)$$

Here, $m = 0, \pm 1, \pm 2$ correspond to the scalar, vector and tensor types. In this paper, we only consider $m = 0$ modes, because they dominate other polarisation modes at the scales we are interested in.

The CMB polarisation components, $E_\ell^{(0)}$ and $B_\ell^{(0)}$, are obtained from the Boltzmann equations for CMB. According to those, $B_\ell^{(0)} = 0$ for all ℓ s and the evolution of $E_\ell^{(0)}$ is given by the following integral solution,

$$\frac{E_\ell^{(0)}(\eta_0, k)}{2\ell+1} = -\frac{3}{2} \sqrt{\frac{(\ell+2)!}{(\ell-2)!}} \int_0^{\eta_0} d\eta \dot{\tau} e^{-\tau} P^{(0)} \frac{j_\ell(k(\eta_0 - \eta))}{(k(\eta_0 - \eta))^2}, \quad (2)$$

where $P^{(0)}$ is the $m = 0$ source term due to Thomson scattering, $\dot{\tau}$ is the differential optical depth for Thomson scattering in conformal time, given by $\dot{\tau} = n_e \sigma_T a$ with the electron number density n_e , the cross section of Thomson scattering σ_T and the scale factor a normalised to the present epoch, and τ is the optical depth between the conformal times η and η_0 . The source term $P^{(0)}$ is related to the initial gravitational potential Φ_0 via the transfer function $D_E(k, \eta)$,

$$P^{(0)} = D_E(k, \eta) \Phi_0. \quad (3)$$

We provide the detailed calculation of the transfer function in the appendix.

We also define the E -modes in the direction $\hat{\mathbf{n}}$ on the sky by

$$E(\hat{\mathbf{n}}) = \sum_{\ell m} (-i)^\ell \sqrt{\frac{4\pi}{2\ell+1}} \int \frac{d^3 k}{(2\pi)^3} E_\ell^{(0)} Y_\ell^m(\hat{\mathbf{n}}). \quad (4)$$

¹ <http://web.haystack.mit.edu/array/MWA>

² <http://www.lofar.org>

³ <http://www.skatelescope.org>

2.2 21 cm line fluctuations

The observed brightness temperature of the 21 cm lines in a direction $\hat{\mathbf{n}}$ and at a frequency ν is given as in Madau et al. (1997) by

$$T_{21}(\hat{\mathbf{n}}; \nu) = \frac{\tau_{21}}{(1 + z_{\text{obs}})} (T_s - T_{\text{CMB}})(\eta_{\text{obs}}, \hat{\mathbf{n}}(\eta_0 - \eta_{\text{obs}})). \quad (5)$$

Here, the conformal time η_{obs} refers to the redshift z_{obs} that satisfies $\nu = \nu_{21}/(1 + z_{\text{obs}})$ with ν_{21} being the frequency corresponding to the 21 cm wavelength. The optical depth for the 21 cm line absorption is

$$\tau_{21} \sim 8.6 \times 10^{-3} x_{\text{H}} \frac{T_{\text{cmb}}}{T_s} \left(\frac{\Omega_{\text{b}} h^2}{0.02} \right) \left[\left(\frac{0.15}{\Omega_{\text{m}} h^2} \right) \left(\frac{1 + z_{\text{obs}}}{10} \right) \right]^{1/2}, \quad (6)$$

where x_{H} is the fraction of neutral hydrogen, and T_s is the spin temperature given by the ratio of the number density of hydrogen atoms in the excited state to that of hydrogens in the ground state.

Equations (5) and (6) show that the observed brightness temperature of the 21 cm lines will reflect the baryon density fluctuations δ_{b} and the fluctuations of the ionised fraction δ_{x} . We can rewrite Eq. (5) in the linear approximation for both δ_{b} and δ_{x} as

$$T_{21}(\hat{\mathbf{n}}; \nu) = [1 - \bar{x}_e(1 + \delta_x)](1 + \delta_b)T_0, \quad (7)$$

where we $x_e = 1 - x_{\text{H}}$ is the ionised fraction and

$$T_0 = 23 \left(\frac{\Omega_{\text{b}} h^2}{0.02} \right) \left[\left(\frac{0.15}{\Omega_{\text{m}} h^2} \right) \left(\frac{1 + z_{\text{obs}}}{10} \right) \right]^{1/2} \text{ mK}. \quad (8)$$

Here, we assume that T_s is much larger than the CMB temperature which is valid soon after the beginning of the reionisation (Ciardi & Madau 2003).

After performing the Fourier transform of Eq. (7), we obtain the 21 cm line fluctuations,

$$\begin{aligned} \delta T_{21}(\hat{\mathbf{n}}; \nu) &\equiv T_{21}(\hat{\mathbf{n}}; \nu) - T_0 \\ &= T_0 \int \frac{dk^3}{(2\pi)^3} [(1 - \bar{x}_e)\delta_{\text{b}} - \bar{x}_e\delta_{\text{x}}] \exp(-i\mathbf{k} \cdot ((\eta - \eta_{\text{obs}})\hat{\mathbf{n}})). \end{aligned} \quad (9)$$

Therefore, the 21 cm line fluctuation map at the frequency ν can be described by

$$\delta T_{21}(\hat{\mathbf{n}}; \nu) = T_0 \sum_{\ell} \int \frac{dk^3}{(2\pi)^3} \sqrt{4\pi(2\ell + 1)} [(1 - \bar{x}_e)(1 + F\mu^2)\delta_{\text{b}} - \bar{x}_e\delta_{\text{x}}] j_{\ell}(k(\eta_0 - \eta_{\text{obs}})) Y_{\ell}^0(\hat{\mathbf{n}}). \quad (10)$$

where we applied Rayleigh's formula to Eq. (9),

$$\exp(-i\mathbf{k} \cdot \mathbf{x}) = \sum_{\ell} \sqrt{4\pi(2\ell + 1)} (-i)^{\ell} j_{\ell}(kr) Y_{\ell}^0(\hat{\mathbf{n}}), \quad (11)$$

where $\hat{\mathbf{n}}$ is measured in the coordinate system where \mathbf{k} is $\hat{\mathbf{e}}_3$. In Eq. (10), we included the factor $(1 + F\mu^2)$ to account for the enhancement of the fluctuation amplitude due to the redshift distortion (Kaiser effect (Kaiser 1987)) on the 21 cm line fluctuations where $\mu = \mathbf{k} \cdot \hat{\mathbf{n}}$ and $F = d \ln g / d \ln a$ with $g(a)$ the linear growth factor of baryon fluctuations (Bharadwaj & Ali 2004).

2.3 Cross-correlation between 21 cm and E -mode fluctuations

The angular power spectrum is defined as the average of the spherical harmonic coefficients $a_{\ell m}$ over the $(2\ell + 1)$ m -values,

$$C_{\ell} = \sum_m \frac{\langle |a_{\ell m}|^2 \rangle}{(2\ell + 1)}, \quad (12)$$

where the $a_{\ell m}$ are defined for an arbitrary sky map $f(\hat{\mathbf{n}})$ as

$$f(\hat{\mathbf{n}}) = \sum_{\ell m} a_{\ell m} Y_{\ell}^m. \quad (13)$$

From Eqs. (4) and (10), we can thus derive the angular power spectrum of the cross-correlation between E -modes and 21 cm line fluctuations

$$\begin{aligned} C_{\ell}^{E-21} &= 4\pi T_0 \int d\eta \int \frac{dk^3}{(2\pi)^3} \frac{\langle E_{\ell}^0(\eta, k) [(1 - \bar{x}_e)(1 + F\mu^2)\delta_{\text{b}} - \bar{x}_e\delta_{\text{x}}] \rangle}{(2\ell + 1)} j_{\ell}(k(\eta_0 - \eta)) \\ &= -\frac{3}{\pi} T_0 \sqrt{\frac{(\ell + 2)!}{(\ell - 2)!}} \int dk \int d\eta k^2 \dot{\tau} e^{-\tau} D_E(k, \eta) \left[\frac{4}{3} (1 - \bar{x}_e) P_{\Phi\delta_{\text{b}}} - \bar{x}_e P_{\Phi\delta_{\text{x}}} \right] \frac{j_{\ell}(k(\eta_0 - \eta_{\text{obs}})) j_{\ell}(k(\eta_0 - \eta))}{(k(\eta_0 - \eta))^2}, \end{aligned} \quad (14)$$

where $P_{\Phi\delta_b}$ and $P_{\Phi\delta_x}$ are the power spectra of the cross-correlation between the gravitational potential and the baryon density fluctuations, and between the gravitational potential and the fluctuations of the ionised fraction, respectively. In the second line, we use $F\langle\mu^2\rangle = 1/3$ for the matter dominated epoch.

The cosmological linear perturbation theory provides the relation between the baryon density fluctuations and the initial gravitational potential (e.g. Kodama & Sasaki 1984), $\delta_b(k, \eta) = k^2 D_b(k, \eta) \Phi_0(k)$ where D_b is the transfer function of baryons. The power spectrum $P_{\Phi\delta_b}$ can thus be written in terms of the initial power spectrum of the gravitational potential P_Φ as

$$P_{\Phi\delta_b} = k^2 D_b(k, \eta) P_\Phi. \quad (15)$$

Although the reionisation process is not well-known, we can reasonably expect that there is a relation between the fluctuations of the ionised fraction and that of the matter density. Ionising sources are formed in dense regions. They ionise the surrounding medium with an efficiency that depends on the density of the medium. This translates into two possible cases: One where ionised fluctuations and matter over-densities coincide, and one where ionised fluctuations and matter density are anti-biased (e.g. Benson et al. 2001). Following Alvarez et al. (2006), we assume that the fluctuations of the ionised rate are associated with the matter density contrast using the Press-Schechter description (Press & Schechter 1974). As a result, the power spectrum $P_{\Phi\delta_x}$ is given by

$$\bar{x}_e P_{\Phi\delta_x} = -\bar{x}_H \ln \bar{x}_H [\bar{b}_h - 1 - f] D_m(k, \eta) k^2 P_\Phi, \quad (16)$$

where D_m is the transfer function of matter (both dark and baryonic), \bar{b}_h is the average bias of dark matter halos more massive than M_{\min} , the minimum mass of the source of ionising photons

$$\bar{b}_h = 1 + \sqrt{\frac{2}{\pi}} \frac{e^{-\delta_c^2/2\sigma^2(M_{\min})}}{f_{\text{coll}}\sigma(M_{\min})}, \quad (17)$$

where $\sigma(M)$ is the variance of the density fluctuations smoothed with a top-hat filter of the scale corresponding to a mass M , and f_{coll} is the fraction of matter collapsed into halos with $M > M_{\min}$. In this paper, we choose M_{\min} such that the halo virial temperature is $T_{\text{vir}}(M_{\min}) = 10^4$ K. This choice corresponds to the assumption that the ionising sources form in dark matter halos where the gas cools efficiently via atomic cooling. The parameter f describes the reionisation regime we are interested in. For $f = 0$, we are in the ‘‘photon-counting limit’’ case where recombinations are not important and where the progress of the reionisation depends on the number of ionising photons only. The over-dense regions contain more collapsed objects which are sources of ionising photons. Therefore, in this case, ionisation in over-dense regions is easier than in under-dense regions. On the contrary, $f = 1$ indicates the ‘‘Strömgren limit’’ case where ionisation is balanced by recombination. Although the over-dense regions contain more sources of ionising photons, the recombination rate in over-dense regions is higher than in under-dense regions. Hence, over-dense regions in the $f = 1$ case have a lower ionised fraction than in the $f = 0$ case (for details, see Alvarez et al. 2006).

Finally, in order to calculate Eq. (14), we need the evolution of the mean neutral hydrogen fraction for which we use a simple parameterisation based on two key quantities, the reionisation epoch (defined as the redshift at which the ionised fraction equals 0.5), z_{re} , and the duration of reionisation, Δz ,

$$\bar{x}_H(z) = \frac{1}{1 + \exp[-(z - z_{\text{re}})/\Delta z]}. \quad (18)$$

In the following we will vary both z_{re} and Δz , and investigate their effect on the cross-correlation signal.

3 RESULTS AND DISCUSSIONS

As one can see in Eq. (14), the angular power spectrum C_ℓ^{E-21} of the cross-correlation is a function of several key parameters that encode the prominent details of the reionisation history. Consequently, and as we show here, the global shape of the C_ℓ^{E-21} as well as the amplitude and redshift dependence of its first peak are features that can be traced back to the reionisation characteristics.

First it is worth noting that because of the double integration of the spherical Bessel functions in Eq. (14), the angular power spectrum reflects the source term $P(k_{\text{obs}})$, namely the quadrupole term of the CMB temperature anisotropy, at z_{obs} where k_{obs} satisfies $k_{\text{obs}} = \ell/(\eta_0 - \eta_{\text{obs}})$. Accordingly, the angular power spectrum of the cross-correlation exhibits its first peak at a multipole $\ell < 10$ which corresponds to the angular separation of the quadrupole at z_{obs} . The oscillations at higher ℓ s (> 10) are due to the free streaming of the quadrupole at redshifts higher than z_{obs} .

We first investigate the effect of the duration of reionisation Δz . We present in the left panel of Fig. 1 three cases: One corresponding to a instantaneous reionisation $\Delta z = 0.01$ and two with longer durations ($\Delta z = 1$ and 2). We compute the angular spectra of the cross-correlation between 21 cm line fluctuations and the CMB E -mode fluctuations for those different durations but for the same given epoch of reionisation, $z_{\text{re}} = 10$, an observing redshift, $z_{\text{obs}} = 10$, and in the case of the ‘‘photon-counting limit’’ ($f = 0$). The difference in the duration of reionisation induces both a minor and a major effects




Figure 1. The left panel shows the angular power spectra of the cross-correlation between E -mode polarisation and 21 cm line fluctuations. We set $z_{\text{re}} = 10$, $z_{\text{obs}} = 10$ and $f = 0$ for all curves. The dotted line represents the angular spectrum for $\Delta z = 0.01$. The solid and dashed lines are for $\Delta z = 1$ and $\Delta z = 2$, respectively. For comparison the thin solid line, shows the angular power spectrum of the CMB E -modes for $z_{\text{re}} = 10$ and $\Delta z = 1$. The right panel is the evolution of the visibility function $\tau e^{-\tau}$ in the same reionisation model as the left panel. The dotted, solid and dashed lines are for $\Delta z = 0.01$, $\Delta z = 1$ and $\Delta z = 2$, respectively.




Figure 2. The angular power spectra of the cross-correlation between E -mode polarisation and 21 cm line fluctuations. The dotted line is for $z = 12$. The solid and dashed lines are for $z = 10$ and $z = 8$, respectively. We use $\Delta z = 1$, $z_{\text{obs}} = 10$ and $f = 0$ for all curves.

on the spectra. The minor effect is the observed shift of the peak position to higher ℓ s with increasing durations. Since a longer duration increases the contribution of the visibility function $\tau e^{-\tau}$ in Eq. (14), which we plot in the right panel of Fig. 1, the contributions from redshifts larger than z_{obs} accumulate in the calculation of the integration. However, spherical Bessel functions have a steep cutoff after η_{obs} so that the contributions from epochs earlier than η_{obs} are negligible. Therefore, additional effective contributions come from only the former epoch $\eta < \eta_{\text{obs}}$, and make the peak shift toward higher ℓ modes.

The major effect of Δz exhibited in the plot (left panel of Fig. 1) is the damping at high ℓ s with increasing durations of reionisation. This occurs again due to the increase of the amplitude of the visibility function at $z > z_{\text{re}}$. At high ℓ s, the source term $P^{(0)}$ exhibits rapid oscillations. Since the tail of the visibility function is larger, the range of the integral is larger, and it thus encompasses more oscillations. As a result, many more cancellations occur which damp the overall power. We note that the damping depends on the duration of the reionisation which is an advantage of observing the cross-correlation signal. As a matter of fact, it is difficult to see differences among CMB polarisation spectra for different durations, because the Thomson optical depth is not strongly sensitive to this parameter. However the plots in the left panel of Fig. 1, clearly show that the damping of angular spectrum of the cross-correlation is sensitive to the duration of reionisation making it a potential tool for constraining this parameter of the reionisation history.

We now vary the reionisation redshift z_{re} keeping the other parameters fixed, and setting $z_{\text{obs}} = 10$, $\Delta z = 1$ and $f = 0$. We find that increasing z_{re} shifts the position of the first peak to small scales (Fig. 2). The reason is the same as the one invoked for the shifts observed in the left panel of Fig. 1. When reionisation happens early, the integration of the E -mode polarisation in Eq. (14) takes into account more contribution from higher redshifts. This induces a shift of the peak position to higher ℓ modes.

Figure 3 shows the evolution of the first peak amplitude as a function of z_{obs} , the redshift of the 21 cm line emission, for the three above-mentioned durations of reionisation Δz . Here, we set $z_{\text{re}} = 10$ and $f = 0$. As shown by Alvarez et al. (2006) using Eq. (18), the amplitude of the cross-correlation of the CMB Doppler temperature anisotropies and the 21 cm line fluctuations




Figure 3. The amplitude of the first peak of the cross-correlation signal as a function of redshift z_{obs} . The dotted, solid and dashed lines indicate the amplitudes for $\Delta z = 0.01$, $\Delta z = 1$ and $\Delta z = 2$, respectively. For both plots, $z_{\text{re}} = 10$ and $f = 0$.




Figure 4. The evolution of the first peak amplitude for different z_{re} . The dotted, solid and dashed lines represent the evolution for $z_{\text{re}} = 12$, $z_{\text{re}} = 10$ and $z_{\text{re}} = 8$, respectively. In this figure, we assume $\Delta z = 1$ and $f = 1$.

reaches its maximum value at $z_{\text{re}} = 10$ where the ionised fraction is about half. The same is true in the case of the cross-correlation of 21 cm line fluctuations and E -mode polarisation. The reason for this is that the CMB polarisation is produced in the ionised regions whereas 21 cm line fluctuations are associated with neutral regions. We note that the signal in Fig. 3 is smaller at lower redshifts than at higher redshifts. This is due to the fact that the 21 cm signal vanishes once reionisation is completed, i.e. at redshifts smaller than z_{re} . For higher redshifts ($z_{\text{obs}} > z_{\text{re}}$), the low scattered quadrupole component of the E -modes is suppressed like in the lower redshifts part, but the integration in the E -modes picks up contributions from the reionisation epoch and the E -modes are correlated with a larger contribution of the 21 cm signal. As a result, the amplitude of the cross-correlation does not decline as steeply as for $z_{\text{obs}} < z_{\text{re}}$. The difference between the curves is easily interpreted. In the $\Delta z = 2$ case, full reionisation takes longer to complete than in the $\Delta z = 1$ case. Therefore, neutral regions persist longer, broadening the skewed bell-shape of the curve centered on $z_{\text{obs}} = z_{\text{re}}$. The slight difference in the peak value is due to the cumulative effect of the integral in Eq. (14). This is particularly well illustrated by the case of “instantaneous reionisation” ($\Delta z = 0.01$). It is worth noting that our assumption, $T_s \gg T_{\text{cmb}}$, is formally not valid before the beginning of reionisation. In order to evaluate the cross-correlation signal correctly, we should rather take into account the full evolution of T_s with redshift in the calculation of the 21 cm line fluctuations. However, by the time the ionisation fraction has risen above the per cent level, efficient Compton and X-ray heating will guarantee that the spin temperature be well above T_{cmb} (e.g. Pritchard & Furlanetto 2007).

The evolution of the first peak amplitude for different z_{re} is shown in Fig. 4. As expected, the amplitude reaches its maximum value at each z_{re} . Since the 21 cm optical depth is higher at high z_{obs} , the amplitude of the 21 cm line fluctuations at higher redshifts is higher than that at lower redshifts. Therefore, and as exhibited by the comparison of the amplitudes for the different z_{re} , the maximum value of the first peak cross-correlation amplitude is larger in the early reionisation case than in the late reionisation case. This again is the same evolution that shows up in the power spectrum of the cross-correlation between the CMB temperature anisotropy and the 21 cm line fluctuations (cf. Fig. 2 of Alvarez et al. 2006). The evolution, with the observed redshift, of the shape and the amplitude of the first peak of cross-correlation between the 21 cm line fluctuations and E -mode fluctuations should allow us to constrain both the duration and redshift of reionisation.




Figure 5. The left panel shows the evolution of the first peak amplitude for two extreme values of the parameter f . The solid and dashed lines are for $f = 0$ and $f = 1$, respectively. We set $z_{\text{re}} = 10$ and $\Delta z = 1$ in all plots. The right panel shows the evolution of the first peak amplitude in the case of a double reionisation model. The solid line (left axis) is for the first peak amplitude at given redshift. The dotted line (right axis) indicates the ionisation fraction in the double reionisation model.

Ultimately, the parameter f is not independent of the epoch z_{re} of reionisation and its duration Δz . However, making that dependency explicit would require the very detailed description of the reionisation processes that we are still lacking today. We can nevertheless illustrate how the cross-correlation power spectrum changes for different global reionisation scenarios. The left panel in Fig. 5 shows the variation of the evolution of the first peak amplitude with the parameter f . There is no difference of the peak position in both spectra for the “photon counting limit” $f = 0$ and the “Strömgren limit” $f = 1$ but the amplitudes are different. When f increases, recombinations become more and more important. Accordingly, over-dense regions in the case of $f = 0$ have higher ionisation fraction than in the case of $f = 1$ as mentioned in Sec. 2.3. Since the fluctuations of the neutral fraction are defined by $\delta_{\text{H}} = -\delta_x$, the absolute value of the 21 cm line fluctuations for $f = 0$ is higher than for $f = 1$. Therefore, the absolute value of the amplitude of the CMB E -mode anisotropies and the 21 cm line fluctuations correlation for $f = 0$ is higher than for $f = 1$ as shown in the left panel of Fig. 5.

Finally, we also calculate the cross-correlation between the 21 cm line fluctuations and the E -mode CMB anisotropies in the case of a double reionisation. For illustrative purposes, we chose a simple parametrisation of the mean ionised fraction

$$\bar{x}_e(z) = 1 - \frac{1}{1 + \exp[-(z - z_{\text{re}})/\Delta z]} + A \exp\left[-\frac{(z - z_{\text{first}})^2}{\Delta z_{\text{first}}^2}\right], \quad (19)$$

which is the simple reionisation case formula with $z_{\text{re}} = 8$ and $\Delta z = 1$ on top of which we added a Gaussian with $A = 0.8$, $z_{\text{first}} = 12$, and $\Delta z_{\text{first}}^2 = 2$ that encodes a first passage through an eighty per cent reionised universe at a redshift of twelve. These parameters have been adjusted so as to reproduce the same value of the optical depth ($\tau = 0.08$) as the single reionisation model shown in the left panel of Fig. 5. The redshift evolution of \bar{x} in this toy model is represented by the dashed line in the right panel of Fig. 5 (read right axis for values of \bar{x}_e). In the same figure, the solid line shows the corresponding evolution of the first peak of the cross-correlation (left axis). Every time the ionised fraction takes the value 0.5 there is a peak in the cross-correlation signal. In the case of our model for double reionisation, we observe three peaks, i.e. the ionisation fraction crosses the 0.5 line thrice. However, the two peaks at $z_{\text{obs}} \sim 8$ and 11 are very close to each other. If two consecutive crossings of the 0.5 line occur very soon after each other, the corresponding peaks in the cross-correlation signal will be essentially indistinguishable. Similarly to what has been found by Alvarez et al. (2006), this behaviour is a very interesting property as it implies that we can use the “trajectory” of the evolution of the peak amplitude to constrain the reionisation model.

4 CONCLUSION

We have examined the cross-correlation between the E -mode CMB polarisation and the fluctuations of the 21 cm line background. In particular, we calculated the angular power spectrum of the cross-correlation on large scales using a simple parametrisation of reionisation, focusing explicitly on the redshift z_{re} of its occurrence and on its duration Δz . We also considered the case of double reionisation for which we obtained specific signatures.

The angular power spectrum of the cross-correlation traces the quadrupole component of the CMB temperature anisotropies at the redshift where the 21 cm line is observed. Both the amplitude and the position of the first peak depend on the reionisation epoch, as well as on the redshift z_{obs} probed by the 21 cm observations. We showed that the signal peaks when z_{obs} matches z_{re} , in the very same way as does the cross-correlation between the CMB Doppler temperature anisotropies and the fluctuations of the 21 cm line background (see Alvarez et al. 2006), that is when the 21 cm observations probe the epoch at which the ionised fraction becomes one half. When probing higher redshifts, the polarisation and 21 cm cross-correlation signal decreases slowly, whereas it falls off rapidly when lower redshifts z_{obs} are considered.

The location of the first peak in the $(z_{\text{obs}} - C_{\ell}^{E-21})$ plane is essentially independent of Δz . Therefore, tracking the evolution of the first peak amplitude by scanning the sky in z_{obs} bins would allow to determine z_{re} . Moreover, the width of the bell-shaped curve describing the evolution with z_{obs} of the C_{ℓ}^{E-21} first peak depends on Δz . The duration of the reionisation is thus also accessible through this observable quantity.

We also showed that, in addition to the amplitude and the position of the first peak, the duration of reionisation induces a significant damping of the oscillations on small scales. This strong dependence on the duration shows specifically in the cross-correlation power spectrum as it does not arise in the auto-correlations power spectra.

In the present study, we assumed the observation of the 21 cm line ideal. In practice, it amounts to consider that the frequency window function is the delta function $\delta(\nu - \nu_{\text{obs}})$, implying that observations are done at the specified frequency ν_{obs} only. Of course, for real observations, the finite instrumental frequency width has to be taken into account. For too broad window functions, this is susceptible to induce an additional damping of the higher ℓ oscillations. We thus checked this effect in the case of the LOFAR interferometer that has a frequency width of 4 MHz. As a result, we did not obtain any additional significant damping of the oscillations for $\ell < 200$.

Finally, due to the cosmic variance that limits CMB measurements on large scales, it may actually be difficult to pin-point the first peak with great accuracy. Moreover, ground-based telescopes designed for observations of the 21 cm line background cannot cover the whole sky. Those two facts are not important limitations for our study, since the damping of oscillations in the cross-correlation power spectrum on intermediate scales ($10 < \ell < 100$), which are well enough sampled, depends strongly on the reionisation evolution. Actually, even if the reionisation duration is so long that we cannot detect the oscillations due to the strong damping, it is still possible to obtain a lower limit for the duration. We will study this in more detail with numerical simulations in the future.

REFERENCES

- Adshead P., Furlanetto S., 2007, ArXiv:0706.3220
 Alvarez M. A., Komatsu E., Doré O., Shapiro P. R., 2006, ApJ, 647, 840
 Barkana R., Loeb A., 2001, Phys. Rep., 349, 125
 Benson A. J., Nusser A., Sugiyama N., Lacey C. G., 2001, MNRAS, 320, 153
 Bharadwaj S., Ali S. S., 2004, MNRAS, 352, 142
 Cen R., 2003, ApJ, 591, 12
 Ciardi B., Madau P., 2003, ApJ, 596, 1
 Cooray A., 2004, Phys. Rev. D, 70, 063509
 Cooray A., 2006, Physical Review Letters, 97, 261301
 Fan X., et al., 2006, AJ, 132, 117
 Field G., 1958, Proc. IREE Aust., 46, 240
 Furlanetto S. R., Oh S. P., Pierpaoli E., 2006, Phys. Rev. D, 74, 103502
 Gunn J. E., Peterson B. A., 1965, ApJ, 142, 1633
 Hu W., Sugiyama N., 1995, ApJ, 444, 489
 Hu W., White M., 1997, Phys. Rev. D, 56, 596
 Jelic V., Zaroubi S., 2008, in preparation
 Kaiser N., 1987, MNRAS, 227, 1
 Kodama H., Sasaki M., 1984, Progress of Theoretical Physics Supplement, 78, 1
 Loeb A., Zaldarriaga M., 2004, Physical Review Letters, 92, 211301
 Madau P., Meiksin A., Rees M. J., 1997, ApJ, 475, 429
 Page L., et al., 2007, ApJS, 170, 335
 Press W.H., Schechter P., 1974, ApJ, 187, 425
 Pritchard J. R., Furlanetto S. R., 2007, MNRAS, 376, 1680
 Salvaterra R., Ciardi B., Ferrara A., Baccigalupi C., 2005, MNRAS, 360, 1063
 Slosar A., Cooray A., Silk J. I., 2007, MNRAS, 377, 168
 Spergel D. N., et al., 2007, ApJS, 170, 377
 Tashiro H., Sugiyama N., 2006, MNRAS, 372, 1060
 Tozzi P., Madau P., Meiksin A., Rees M. J., 2000, ApJ, 528, 597
 Wouthusen S., 1952, AJ, 57, 31
 Zaldarriaga M., 1997, Phys. Rev. D, 55, 1822

APPENDIX A: THE CMB E -MODE POLARISATION

The source term of E -mode polarisation in Eq. (2) (Hu & White 1997),

$$P^{(0)}(\eta, k) = \Theta_2^{(0)}(\eta, k) - \sqrt{6}E_2^{(0)}(\eta, k), \quad (\text{A1})$$

where $\Theta_2^{(0)}$ is the quadrupole moment of the CMB temperature anisotropies.

The CMB temperature anisotropies are evaluated by the following integral solutions obtained from the Boltzmann equations,

$$\frac{\Theta_\ell^{(0)}(\eta_0, k)}{2\ell + 1} = \int_0^{\eta_0} d\eta e^{-\tau} \sum_{\ell'} S_{\ell'}^{(0)}(\eta) j_\ell^{(\ell')} (k(\eta_0 - \eta)), \quad (\text{A2})$$

where the $j_\ell^{(\ell')}$ are expressed in terms of the spherical Bessel functions $j_\ell(x)$,

$$j_\ell^{(0)}(x) = j_\ell(x), \quad j_\ell^{(1)}(x) = j_\ell'(x), \quad j_\ell^{(2)}(x) = \frac{1}{2}[3j_\ell''(x) + j_\ell(x)]. \quad (\text{A3})$$

and $S_\ell^{(0)}$ are source terms written as

$$S_0^{(0)} = \dot{\tau}\Theta_0^{(0)} - \dot{\Phi}, \quad S_1^{(0)} = \dot{\tau}v_B + k\Psi, \quad S_2^{(0)} = \frac{\dot{\tau}}{10} [\Theta_2^{(0)} - \sqrt{6}E_2^{(0)}], \quad (\text{A4})$$

where v_b is the velocity of the baryon fluid, Φ and Ψ are the scalar-type metric perturbations in the conformal Newtonian gauge. The cosmological linear perturbation theory provides the evolution of Φ and Ψ and the initial condition of Θ_0 in the matter dominated epoch as (Kodama & Sasaki 1984)

$$\Psi = -\Phi = -\frac{9}{10}\Psi(0), \quad \Theta_0(0) = \frac{3}{5}\Psi(0), \quad (\text{A5})$$

where $\Psi(0)$ is the initial metric perturbation.

Before recombination, photons and baryons are tightly coupled by Compton scattering. Hence, the tight-coupling approximation is a strong tool for solving Eqs. (2) and (A2). Hu & Sugiyama (1995) provided a useful analytic expression of the monopole component in this approximation,

$$\Theta_0(\eta) + \Phi(\eta) = [\Theta_0(0) + \Phi(0)] \cos kr_s(\eta) - \frac{k}{\sqrt{3}} \int_0^\eta d\eta' [\Phi(\eta') - \Psi(\eta')] \sin[kr_s(\eta) - kr_s(\eta')], \quad (\text{A6})$$

where r_s is the integration of the sound speed $c_s = 1/\sqrt{3(1+R)}$,

$$r_s = \int d\eta \frac{1}{\sqrt{3(1+R)}}. \quad (\text{A7})$$

The dipole and quadrupole components, and the baryon velocity are written as

$$\Theta_1 = -\frac{3}{k}(\dot{\Theta}_0 + \dot{\Phi}), \quad \Theta_2 = \frac{4\sqrt{4}}{9} \frac{k}{\dot{\tau}} \Theta_1, \quad v_B = \Theta_1. \quad (\text{A8})$$

Accordingly, the quadrupole component of E -modes is given by

$$E_2 = -\frac{\sqrt{6}}{4} \Theta_2. \quad (\text{A9})$$

After recombination, the tight-coupling approximation is no longer valid. Using Eqs. (A5)-(A9) as initial conditions, we solve Eqs. (2) and (A2) numerically in order to obtain the transfer function $D_E(k, \eta)$ in Eq. (3).