LSS2014 PRESENTATION TOPICS

Dear Cosmic Structure Formation 2014 students,

In this document you find a listing of suggested topics for the final presentation. Mentioned with most topics are "some" literature suggestions. In some cases this concerns an essential key reference which then would advisably be used for the report. In other cases you find a listing of publications that would provide you with a good overview of important papers relating to the subject, and it will be up to you to select the ones you need (in addition of course to other works you find yourself as revelant).

Take care that your presentation is a review of relevant work, not JUST a summary of one or two papers. It should be a critical discussion, and place the topic within the context of ongoing reseach.

Have fun, looking forward to your presentations, Rien van de Weygaert Kapteyn Institute, 9700 AV Groningen tel.: 050-3634086 e-mail: weygaert@astro.rug.nl

1. Evidence for Dark Energy.

- A discussion on the observational evidence for a cosmological constant Lambda or, in a wider context, for the reality of Dark Energy. Is it realistic to measure the real equation of state of the vacuum ?
- Literature suggestions:
 - Albrecht A., Bernstein G., Cahn R., Freedman W., et al., 2006 Report of the Dark Energy Task Force astro-ph
 - Carroll S.M., Press W.M., 1992 The Cosmological Constant ARAA 30, 499
 - Huterer D., Turner M.S., 2001
 Probing dark energy: Methods and strategies
 Phys. Rev. D., 64, 123527
 - Kirshner R.P, 2003 Throwing Light on Dark Energy Science 300, 1914
 - Padmanabhan T., 2003
 Cosmological Constant the Weight of the Vacuum hep-th/0212290
 - Riess A.G., Turner M.S., 2004
 From Slowdown to Speedup
 Sci. Am., feb. 2004, pg. 62
 - Trotta R., Bower R., 2006
 Surveying the Dark Side
 Astronomy. Geophys. 47, 422
 - Weinberg S., 1989
 The cosmological constant problem
 Rev. Mod. Phys. 61, 1

2. Baryonic Oscillations

- Baryonic oscillations in the power spectrum of galaxy clustering have recently been detected from the 2dFGRS and SDSS galaxy redshift survey. At the moment various deep redshift surveys are being designed to measure the baryonic oscillations and get a precise measurement of the equation of state of dark energy from this purely geometric and robust method. Give a discussion of the fundamental physics behind the oscillations, on the detection and measurement of the oscillations and on the prospects to study dark energy.

- Literature suggestions:

- Cole S., et al., 2005
 The 2dF Galaxy Redshift Survey: power-spectrum analysis of the final data set and cosmological implications
 MNRAS, 362, 505
- Eisenstein D.J., Hu W., 1997
 Baryonic Features in the Matter Transfer Function
 ApJ 496, 605
- Eisenstein D.J., et al., 2005
 Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies
 ApJ, 633, 560
- Glazebrook K., et al., 2007 The WiggleZ project: AAOmega and Dark Energy ASP, astro-ph/0701876
- Parkinson D., et al., 2007 Optimizing Baryon Acoustic Oscillation Surveys - I. Testing the concordance LCDM cosmology

MNRAS, astro-ph/0702040

- Percival et al., 2006
 Measuring the Matter Density using Baryonic Oscillations in the SDSS ApJ, astro-ph/0608635
- Seo H-J, Eisenstein D.J., 2005
 Baryon Acoustic Oscillations in Simulated Galaxy Redshift Surveys
 ApJ 633, 575
- Seo H-J, Eisenstein D.J., 2005
 Probing Dark Energy with Baryonic Acoustic Oscillations ASP, Vol. 399

 Seo H-J, Eisenstein D.J., 2007 Improved forecasts for the baryon acoustic oscillations and cosmological distance scale astro-ph/0701079
 White M 2005

White M., 2005
 Baryon Oscillations
 Astroparticle Physics 24, 334

3. The Cosmic Web

- On scales of a few up to more than a hundred Megaparsec displays galaxies delineate a salient and pervasive weblike pattern, the *Cosmic Web* (Bond et al. 1996). Galaxies and mass are arranged into elongated filaments, sheetlike walls and dense compact clusters. These surround large *voids*, enormous regions with sizes in the range of $20-50h^{-1}$ Mpc that are practically devoid of any galaxy, usually roundish in shape and occupying the major share of space in the Universe. Write a review on observational aspects of the cosmic web, along with a discussion of N-body simulations revealing the same weblike structure. Literature is too vast to provide any decent measure of fitting references. Please contact RvdW. A few suggestions ...
- Literature suggestions:
 - Bond J.R., Kofman L., Pogosyan D., 1996
 How filaments of galaxies are woven into the cosmic web Nature, 380, 603
 - Dolag K. et al., 2005
 Simulating the physical properties of dark matter and gas inside the cosmic web MNRAS, 2005/2006, astro-ph/0511357
 - Gott J.R. III, et al., 2005
 A map of the Universe
 ApJ 624, 463
 - Lahav O., Suto Y., 2003 Measuring our universe from galaxy redshift surveys review, astro-ph/0310642
 - Springel V., Frenk C.S., White S.D.M., 2006 The large-scale structure of the Universe review, Nature, 440, 1137
 - van de Weygaert R., 2002
 Froth across the Universe
 invited review, Modern Theoretical and Observational Cosmology, eds. Plionis & Costakis, Kluwer
 contact RvdW
 - van de Weygaert R., Schaap W., 2007
 the Cosmic Web: Geometrical Analysis and Modelling
 lecture, summerschool Cosmological Data Analysis, Valencia, 2004
 contact RvdW (only concers first part).

4. Voids and the Large Scale Structure

- Voids are a major component of the Megaparsec Universe. With the arrival of the 2dFGRS and SDSS galaxy redshift surveys it is finally possible to obtain a global picture of their characteristics. Review their properties, discuss their role within the formation of large scale structure (both from an observational and theoretical point of view), discuss their sensitivity to cosmological parameters and evaluate what void galaxies can teach about galaxy formation.
- Literature suggestions:
 - Furlanetto S.R., Piran T., 2006
 The evidence of absence: galaxy voids in the excursion set formalism MNRAS 366, 467
 - Hoyle F., Vogeley M.S., 2004
 Voids in the Two-degree Field Galaxy Redshift Survey
 ApJ 607, 751
 - Kirshner R., Oemler A., Schechter P.L., Shectman S.A., 1987
 A survey of the Bootes Void
 ApJ 314, 493
 - Park D., Lee J., 2006
 The Void Ellipticity Distribution as a Probe of Cosmology astro-ph/0610520
 - Peebles P.J.E., 2001
 The Void Phenomenon
 ApJ 557, 495
 - Rojas R.R., Vogeley M.S., Hoyle F., Brinkmann J., 2004
 Photometric properties of void galaxies in the Sloan Digital Sky Survey
 ApJ 617, 50
 - Ryden B., Melott A.L., 1996
 Voids in Real Space and in Redshift Space
 ApJ, 470, 160
 - Shandarin S., Feldman H.A., Heitmann K., Habib S., 2005 Shapes and Sizes of Voids in the LCDM Universe: Excursion Set Approach
 - Sheth R., van de Weygaert R., 2004
 A hierarchy of voids: much ado about nothing MNRAS 350, 517

 van de Weygaert R., van Kampen E., 1993
 Voids in gravitational instability scenarios - I. Global density and velocity field in an Einstein-de Sitter universe
 MNRAS 263, 481

5. Hierarchical Clustering and the Excursion Set Formalism

- Structure in the Universe builds up in a hierarchical fashion: the first objects to form are small, larger ones form by the gradual merging of smaller clumps. To see how such structures emerge out of an initially Gaussian density field a sophisticated mathematical formalism has been developed, the socalled *excursion set formalism*. This formalism allows the computation of the resulting mass spectrum of galaxies and clusters, the development of the void distribution, even that of the *Cosmic Web* and the merging history of halos in a hierarchically evolving mass distribution. Discuss the essential ingredients of this elegant formalism, establish the link with diffusion equations, and evaluate its successes and limitations.
- Literature suggestions:
 - Appel L., Jones B.J.T., 1990
 The mass function in biased galaxy formation scenarios
 MNRAS 245, 522
 - Bond J. R., Cole S., Efstathiou G., Kaiser N.
 Excursion set mass functions for hierarchical Gaussian fluctuations ApJ., 379, 440
 - Bower R.G., 1991
 The evolution of groups of galaxies in the Press-Schechter formalism MNRAS 248, 322
 - Hanami H., 2001 Statistics of merging peaks of random Gaussian fluctuations: skeleton tree formalism

MNRAS 327, 721

- Lacey C., Cole S., 1993
 Merger rates in hierarchical models of galaxy formation MNRAS 262, 627
- Lacey C., Cole S., 1994
 Merger rates in hierarchical models of galaxy formation II. Comparison with N-body simulations
 MNRAS 271, 676
- Peacock J.A., Heavens A.F, 1990
 Alternatives to the Press-Schechter cosmological mass function MNRAS, 243, 133

Press W., Schechter P., 1974
Formation of Galaxies and Clusters of Galaxies by Self-Similar Gravitational Condensation
ApJ., 187, 425
Shen J., Abel T., Mo H.J., Sheth R.K., 2006
An Excursion Set Model of the Cosmic Web: the abundance of sheets filaments

An Excursion Set Model of the Cosmic Web: the abundance of sheets, filaments, and halos

ApJ 645, 783

 Sheth R., 1998
 An excursion set model for the distribution of dark matter and dark matter haloes MNRAS 300, 1057

 Sheth R., Mo H.J., Tormen G., 2001
 Ellipsoidal collapse and an improved model for the number and spatial distribution of dark matter haloes
 MNRAS 323, 1

- Sheth R., Tormen G., 2002

An excursion set model of hierarchical clustering: ellipsoidal collapse and the moving barrier $% \left(\mathcal{A}^{\prime}\right) =\left(\mathcal{A}^{\prime}\right) \left(\mathcal{A}^{\prime}\right)$

MNRAS, 329, 61

 Sheth R., van de Weygaert R., 2004
 A hierarchy of voids: much ado about nothing MNRAS 350, 517

 $\rm astro-ph/0611454$

Zentner A.R., 2006
 The Excursion Set Theory of Halo Mass Functions, Halo Clustering, and Halo Growth
 active ph/0611454

6. Cluster Evolution and Intracluster Medium

- Clusters of galaxies are the most massive fully collapsed and virialized objects in the Universe. In fact, they are huge balls of hot $10^7 10^8$ K intracluster gas. Describe the evolution of clusters, focussing in particular on that of the hot X-ray emitting gas.
- Literature suggestions:
 - Arnaud M., 2005 X-ray observations of Clusters of Galaxies Societa Italiana di Fisica, astro-ph/0508159 - Borgani S., Guzzo L., 2001 X-ray clusters of galaxies as tracers of structure in the Universe Nature 409, 39 Cohn J.D., White M., 2005 The formation histories of galaxy clusters Astroparticle Physics 24, 316 - Fabian A.C., 1994 Cooling Flows in Clusters of Galaxies ARAA, 32, 277 - Kaiser N., 1991 Evolution of Clusters of Galaxies ApJ, 383, 104 - Voit G.M., 2005 Tracing cosmic evolution with clusters of galaxies Rev. Mod. Phys. 77, 207
 - Markevitch M., Vikhlinin A., 2007
 Shocks and Cold Fronts in Galaxy Clusters
 Physics Reports, 67 pp, astro-ph/0701821

7. Clusters and the Sunyaev-Zel'dovich effect

- The hot intracluster gas in clusters may scatter CMB photons to higher frequencies. The resulting spectrum is no longer blackbody. This effect, the Sunyaev-Zel'dovich effect, may open the way towards mapping all clusters throughout the observable Universe. There are two effects, the thermal and the kinetic SZ effect. Describe the effect, present some recent results and analyze the prospects for future experiments like the Planck satellite.

- Literature suggestions:

- Birkinshaw M., 1999
 The Sunyaev-Zel'dovich effect
 Physics Reports 310, 97
- Carlstrom J.E., Holder G.P., Reese E.D., 2002 Cosmology with the Sunyaev-Zeldovich Effect ARAA 40, 643
- Geisbuesch J., Hobson M.P., 2006 Cosmology with the Planck cluster sample MNRAS
- Schäfer B.M. et al., 2004
 Detecting Sunyaev-Zel'dovich clusters with Planck: I. Construction of all-sky thermal and kinetic SZ-maps
 MNRAS
- Sunyaev R.A., Zeldovich Ya.B., 1980
 Microwave background radiation as a probe of the contemporary structure and history of the universe
 ARAA, 18, 537
- Sunyaev R.A., Zeldovich Ya.B., 1980
 The velocity of clusters of galaxies relative to the microwave background The possibility of its measurement
 MNRAS 190, 413
- Sunyaev R.A., Zeldovich Ya.B., 1970
 The Spectrum of Primordial Radiation, its Distortions and their Significance
 Comments on Astrophys. Space Physics, 2, 66
- Sunyaev R.A., Zeldovich Ya.B., 1972
 The Observations of Relic Radiation as a Test of the Nature of X-Ray Radiation from the Clusters of Galaxies
 Comments Astrophys. Space Phys. 4, 173

Vale C., White M., 2006 Finding clusters in SZ surveys New Astron. 11, 207
Voit G.M., 2005 Tracing cosmic evolution with clusters of galaxies Rev. Mod. Phys. 77, 207

8. Cosmology and Clusters of Galaxies

- Clusters of galaxies are the most massive fully collapsed and virialized objects in the Universe. The population of clusters and its evolution are sensitive probes of underlying cosmology and of the spectrum of density fluctuations. In particular interesting is the mass spectrum n(M) of clusters of mass M. On the basis of theoretical predictions one can infer its sensitivity to the cosmological mass density Ω_m , on the power spectrum of density fluctuations and in particular its amplitude (expressed in terms of σ_8 , the root mean square of density fluctuations within a sphere of radius $8h^{-1}$ Mpc). Discuss the theoretical framework of predicted cluster mass spectra (both analytically as well as by means of N-body simulations). In addition, discuss the constraints on cosmological parameters obtained from such studies, in particular on the basis of samples of X-ray clusters. Finally, show how these in combination with CMB experiments yield estimates of in particular Ω_m and Ω_{Λ} .
- Literature suggestions:
 - Bahcall N., Cen R., 1993
 The Mass Function of Clusters of Galaxies
 Astrophys. J., 407, L49
 - Bond J. R., Cole S., Efstathiou G., Kaiser N.
 Excursion set mass functions for hierarchical Gaussian fluctuations Astrophys. J., 379, 440
 - Borgani S., Guzzo L., 2001 X-ray clusters of galaxies as tracers of structure in the Universe Nature 409, 39 Cohn J.D., White M., 2005

The formation histories of galaxy clusters Astroparticle Physics 24, 316

- Eke V., Cole S., Frenk C., 1996
 Cluster evolution as a diagnostic for Ω
 MNRAS, 282, 263
- Eke V., Cole S., Frenk C., 1998 Measuring Ω_o using cluster evolution MNRAS 298, 1145
- Jenkins A., et al., 2001 The mass function of dark matter halos MNRAS, 321, 372

- Peacock J.A., Heavens A.F, 1990
 Alternatives to the Press-Schechter cosmological mass function MNRAS, 243, 133
- Press W., Schechter P., 1974
 Formation of Galaxies and Clusters of Galaxies by Self-Similar Gravitational Condensation
 Astrophys. J., 187, 425
- Sheth R., 1998
 An excursion set model for the distribution of dark matter and dark matter haloes MNRAS 300, 1057
- Sheth R., Mo H.J., Tormen G., 2001
 Ellipsoidal collapse and an improved model for the number and spatial distribution of dark matter haloes
 MNRAS 323, 1
- Sheth R., Tormen G., 2002
 An excursion set model of hierarchical clustering: ellipsoidal collapse and the moving barrier
 MNRAS, 220, 61

MNRAS, 329, 61

- Pierpaoli E., Scott D., White M., 2001
 Power-spectrum normalization from the local abundance of rich clusters of galaxies MNRAS, 325, 77
- Reiprich T.H., Boehringer H., 2002
 The Mass Function of an X-ray flux-limited sample of galaxy clusters
 Astrophys. J., 567, 716
- Rosati P, Borgani S., Norman C., 2002
 The Evolution of X-ray Clusters of Galaxies
 ARAA, 40, 539-77
- Viana P.T.P., Liddle A.R., 1996
 The cluster abundance in flat and open cosmologies
 MNRAS 281, 323
- White M., 2002 The Mass Function ApJS, 143, 241

9. Warm and Hot Intergalactic Medium (WHIM)

- A significant fraction of the baryons in the present universe is hiding in as shock-heated warm-hot intergalactic medium spread throughout the Cosmic Web. Provide a review on the simulations of the Intergalactic medium and the predictions for tracing this gaseous medium. Also include a critical discussion of the claims for having detected this medium and the prospects for mapping the IGM.

- Literature suggestions:

- Cen R., Ostriker J.P., 1999
 Where are the baryons
 ApJ, 514, 1
- Cen R., Ostriker J.P., 2006
 Where are the baryons II: Feedback effects
 ApJ, 650, 560
- Cen R., Ostriker J.P., 2006
 Where are the baryons III: Nonequilibrium effects and observables
 ApJ, 650, 573
- Davé R., et al.

Baryons in the Warm-Hot Intergalactic Medium ApJ, 552, 473

- Kaastra J., et al. 2002

XMM-Newton confirmation of Soft X-ray excess emission in clusters of galaxies the discovery of O VII emission from an extended warm baryonic component A&A, astro-ph/0210684

- Kravtsov A., Klypin A., Hoffman Y., 2002

Constrained simulations of the real universe II. Observational signatures of intergalactic gas in the Local Supercluster Region

ApJ 571, 563

- Nicastro F., 2002

Chandra discovery of a tree in the X-ray forest towards PKS 2155-304: the Local Filament ?

ApJ 573, 157

- Nicastro F. et al., 2005

The mass of the missing baryons in the X-ray forest of the warm-hot intergalactic medium

Nature, 433, 495

- Viel M., et al., 2003

Detecting X-ray filaments in the low redshift Universe with XEUS and Constellation- \blacksquare X

MNRAS, astro-ph/0210497

 Viel M., et al., 2004
 Tracing the Warm Hot Intergalactic Medium in the Local Universe MNRAS, astro-ph/0412566

10. Cosmic Flows and Redshift Space Distortions

- Due to the matter migration flows accompanying the formation of structure in the Universe, maps of galaxies in redshift space are a combination of their true spatial distribution and systematic distortions resulting from the peculiar velocities of galaxies. Investigate the possible effects, provide an overview on the attempt to theoretically describe this and discuss results on the basis of (recent) galaxy redshift surveys.
- Literature suggestions:
 - Dekel, A., 1994
 Dynamics of Cosmic Flows
 ARAA, 32, 371-418
 - Strauss, M., Willick, J., 1995
 The density and peculiar velocity fields of nearby galaxies
 Phys. Rep. 261, 271-431
 - Hamilton, A.J.S.
 Linear Redshift Distortions: A Review
 Invited review Ringberg Workshop on Large-Scale Structure, 1996
 astro-ph/9708102
 - Szalay, A., Matsubara, T., Landy, S., 1998
 Redshift-Space distortions of the Correlation Function in Wide-Angle Galaxy Surveys
 Arc L 408, L1 L4

ApJ, 498, L1-L4

 Madgwick et al., 2003
 The 2dF Galaxy Redshift Suvery: galaxy clustering per spectral type astro-ph/0303668

11. Weak Lensing by Large Scale Structure

- The finding of cosmic shear induced by the inhomogeneous matter distribution in the Universe has triggered enormous activity in trying to infer cosmological parameters, estimating the cosmic power spectrum and ultimately mapping the dark matter distribution throughout the Universe on the basis of the slight distortions of galaxy images by weak lensing by the cosmic mass distribution. Provide a review of the theoretical predictions, of analysis methods, and achieved results, culminating in the first map of the cosmic dark matter distribution.
- Literature suggestions:
 - Bacon D.J., Taylor A.N., 2003
 Mapping the 3D dark matter potential with weak shear
 MNRAS 344, 1307
 - Heavens A., 2003
 3D weak lensing
 MNRAS 343, 1327
 - Kaiser N., Squires G., 1993
 Mapping the Dark Matter with Weak Gravitational Lensing ApJ, 404, 441
 - Kaiser N., 1998
 Weak Lensing and Cosmology ApJ, 498, 26
 - Jain B., Seljak U., 1997
 Cosmological Model Predictions for Weak Lensing: Linear and Nonlinear Regimes
 ApJ, 484, 560
 - Jain B., Seljak U., White S.D.M., 2000
 Ray-tracing simulations of weak lensing by Large-Scale Structure ApJ., 530, 547-577
 - Massey et al., 2007
 Dark matter maps reveal cosmic scaffolding
 Science 445, 286
 - Mellier Y., 1999
 Probing the Universe with Weak Lensing
 ARAA, 37, 127

- Munshi D., Valageas P., van Waerbeke L., Heavens A., 2006 Cosmology with Weak Lensing Surveys astro-ph/0612667, 98 pp.
- Refregier A., 2003
 Weak Gravitational Lensing by Large-Scale Structure ARAA 41, 645
- Schneider P., 2005
 Weak Gravitational Lensing
 Saas-Fee course 2005, astro-ph/0509252, 180pp.
- Vale C., White M., 2003
 Simulating Weak Lensing by Large Scale Structure
 Astroparticle Physics 22, 19
- Van Waerbeke L., Mellier Y., 2003
 Gravitational lensing by Large Scale Structures: A Review astro-ph/0305089

12. Power Spectrum Estimation

- The density fluctuation power spectrum P(k) is one of the most essential ingredients for any theory of structure formation: to a large extent it determines the mode, nature and rate of the emergence of galaxies, clusters and all elements of the cosmic foam in the Universe. Trying to determine the power spectrum from the many sources of information on cosmic structure is one of the major activities of cosmological studies. Provide a discussion of the various methods. Concentrate in particular on the machinery that has been developed to extract significant estimates of P(k) from galaxy redshift surveys.
- Literature suggestions:
 - Feldman H.A., Kaiser N., Peacock J.A., 1994
 Power-spectrum Analysis of Three-dimensional Redshift Surveys ApJ., 426, 23-37
 - Vogeley M.S., Szalay A.S.
 Eigenmode Analysis of Galaxy Redshift Surveys. I. Theory and Methods ApJ., 465, 34-53
 - Tegmark M., Taylor A.N., Heavens A.F., 1997
 Karhunen-Loève Eigenvalue Problems in Cosmology: How should we tackle large data sets
 ApJ., 480, 22-35
 - Hamilton A.J.S., 1997
 Towards optimal measurement of power spectra I. Minimum variance pair weighting and the Fisher matrix
 MNRAS 289, 285-294
 - Hamilton A.J.S., 1997
 Towards optimal measurement of power spectra II. A basis of positive, compact, statistically orthogonal kernels
 MNRAS 289, 295-304
 - Tegmark M., Hamilton A.J.S., Xu Y., 2002
 The power spectrum of galaxies in the 2dF 100k redshift survey MNRAS 335, 887

13. Cosmic Microwave Background Radiation: Thermalization and Spectral Distortions

- The cosmic microwave background radiation provides the strongest observational foundation for the standard Hot Big Bang theory. The fact that its electromagnetic spectrum is almost perfectly Blackbody sets severe constraints on 1) the thermalization processes and epoch of the radiation and 2) spectrum distorting processes. While the anisotropies in the CMB provide a wealth of information on the Universe around the epoch of recombination ($z \sim 1089$), the CMB Planck spectrum forms a probe of the Universe out to redshifts $z \sim 10^7$. Investigate in detail the physical processes that lead to the thermalization of the CMB, and discuss the possible observational spectral signatures incurred by distorting energetic processes at $z < 10^7$. Discuss the observational constraints on the various observational signatures.
- Literature suggestions:
 - Sunyaev R.A., Zeldovich Ya.B., 1980
 - Microwave background radiation as a probe of the contemporary structure and history of the universe

ARAA, 18, 537

- Stebbins A., 1997

The Cmbr-Spectrum a Theoretical Introduction

The Cosmic Microwave Background, Kluwer Academic Press;

- Ed. C.H. Lineweaver, J.G. Bartlett, A. Blanchard, M. Signore, and J. Silk, p.241
- Hu W., 1995
 - Wandering in the Background: A Cosmic Microwave Background Explorer Ph.D. thesis, Univ. California at Berkeley

website: http://background.uchicago.edu/~whu/thesis/thesispage.html

- Zeldovich Ya.B., Sunyaev R.A., 1969
 The Interaction of Matter and Radiation in a Hot-Model Universe Astrophysics and Space Science, Vol. 4, p.301
- Sunyaev R.A., Zeldovich Ya.B., 1970
 The Interaction of Matter and Radiation in a Hot-Model Universe, II
 Astrophysics and Space Science, Vol. 7, p.20
- Sunyaev R.A., Zeldovich Ya. B., 1970
 The spectrum of primordial radiation, its distortions and their significance
 Comments on Astrophysics and Space Physics, vol.2, p.66-73

- Chan K.L, Jones B.J.T., 1975 Distortions of the $3^{\circ}K$ background radiation spectrum: observational constraints on the early thermal history of the Universe Astrophys. J., 195, 1 - Danese L, De Zotti G., 1977 The relic radiation spectrum and the thermal history of the Universe Nuovo Cimento, Rivista, Serie 2, vol. 7, July-Sept. 1977, p. 277-362. - Danese L., De Zotti G., 1982 Double Compton Process and the Spectrum of the Microwave Background A&A, 107, 39 - Burigana C., Danese L., De Zotti G., 1991 Formation and evolution of early distortions op the microwave background spectrum: a numerical study A&A, 246, 49 - Burigana C., De Zotti G., Danese L., 1991 Constraints on the Thermal History of the Universe from the Cosmic Microwave Background Spectrum A&A, 379, 1 - Hu W., Silk J., 1993 Thermalization and spectral distortions of the cosmic background radiation Phys. Rev. D. 48, 485 - Fixsen et al., 1996 The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set Astrophys. J., 473, 576

14. CMB Experiments: Maps, Power Spectra and Cosmological Parameter Estimation

- On the basis of the raw data of CMB temperature measurements across the sky obtained through microwave background experiments (e.g. COBE, Boomerang and WMAP), a data analysis machinery should process this into maps of the MWB sky, of the power spectrum C_l and ultimately significant estimates of cosmological parameters. Describe the various steps of this process, in this concentrating particularly on the mapmaking procedures and the estimation of the power spectrum. The latter involves a discussion on the statistical and numerical methods developed to obtain optimal estimates of C_l , including a description of the CMBFAST package for fast theoretical power spectrum computation.
- Literature suggestions:
 - Hu W., Dodelson S., 2002
 Cosmic Microwave Background Anisotropies
 ARAA, 40, 171-216
 - Seljak U., Zaldarriaga M., 1996
 Line-of-sight integration approach to cosmic microwave background anisotropies
 Astrophys. J. 469, 437
 - Jungman G., Kamionkowski M., Kosowsky A., Spergel D., 1996 Cosmological Parameter Determination with Microwave Background Maps Phys. Rev. D., 54, 1332
 - Bond J.R., Efstathiou G., Tegmark M., 1997
 Forecasting cosmic parameter errors from microwave background anisotropy experiments
 MNRAS 291, L33
 - Zaldarriaga M., Spergel D., Seljak U., 1997
 Microwave Background Constraints on Cosmological Parameters Astrophys. J., 488, 1
 - Tegmark M., 1997
 CMB mapping experiments: A designer's guide Phys. Rev. D., 56, 4514
 - Tegmark M., 1997
 - How to make maps from cosmic microwave background data without losing information

Astrophys. J., 480, L87

- Bond J.R., Jaffe A.H., Knox L., 1998
 Estimating the power spectrum of the cosmic microwave background Phys. Rev. D., 57, 2117
- Bennett C.L. et al., 2003
 First-year WMAP Observations: Preliminary Maps and Basic Results
 ApJS, 148, 1
- Hinshaw G. et al., 2003
 First-year WMAP Observations: Data processing methods and systematic error limits
 - ApJS, 148, 63
- Hinshaw G., et al., 2003
 First-year WMAP Observations: the Angular Power Spectrum ApJS, 148, 135
- Spergel D.N., et al., 2003
 First-year WMAP Observations: Determination of Cosmological Parameters ApJS, 148, 175
- Verde L., et al., 2003
 First-year WMAP Observations: Parameter Estimation Methodology
 ApJS, 148, 195
- Page L., et al., 2003 First-year WMAP Observations: Interpretation of the TT and TE angular power spectrum peaks

ApJS, 148, 233

- Spergel D.N., et al., 2006 Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology astroph
- M. Tegmark, website: Max Tegmark's CMB data analysis center http://www.hep.upenn.edu/~max/cmb/pipeline.html
- WMAP website: http://map.gsfc.nasa.gov/m_mm/pub_papers/firstyear.html
- Healpix website: http://www.eso.org/science/healpix/
- CMBFAST website: http://cmbfast.org/

- Hu W., website:

Cosmological Parameters in the CMB http://background.uchicago.edu/~whu/metaanim.html

15. Cosmic Microwave Background Anisotropies: Characteristic Angular Scales

- Four characteristic angular scales are imprinted on the spectrum of Microwave Background Anisotropies. These are the scale corresponding to the radiation-matter equality ℓ_{eq} , the scale ℓ_A corresponding to the cosmic sound horizon at the epoch of decoupling, the diffusion scale ℓ_D due to (Silk) scattering diffusion and finally the scale $\ell_{K\Lambda}$ at which either the curvature or the cosmic constant takes over from matter as dominant factor behind the expansion of the Universe. Provide a discussion how the various relevant physical processes (Sachs-Wolfe, acoustic oscillations, Silk diffusion damping, Integrated Sachs-Wolfe, etc.) are relating to these characteristic scales and how this leads to differences in angular power spectrum of cosmic microwave background fluctuations.
- Literature suggestions:
 - Hu W., Dodelson S., 2002
 Cosmic Microwave Background Anisotropies
 ARAA, 40, 171-216
 - Hu W., Sugiyama N., Silk J., The Physics of the Microwave Background Anisotropies Nature, Vol. 386, p. 37-43
 - Hu W., 1995
 - Wandering in the Background: A Cosmic Microwave Background Explorer Ph.D. thesis, Univ. California at Berkeley
 - website: http://background.uchicago.edu/ \sim whu/thesis/thesispage.html
 - Hu W.

Cosmic Microwave Background lecture notes, 172 pp. contact RvdW for pdf file

- Hu W., Sugiyama N., 1995
 Anisotropies in the Cosmic Microwave Background: an Analytic Approach ApJ., 444, 489
- Hu W., White M., 2004
 The Cosmic Symphony
 Sci. Am., feb. 2004, 44

- Bond J.R., 1996

Theory and Observations of the Cosmic Background Radiation
in "Cosmology and Large Scale Structure", Les Houches LX, eds. R. Schaeffer, J. Silk, M. Spiro, J. Zinn-Justin, Elsevier
(contact RvdW for pdf file).
Seljak U., Zaldarriaga M., 1996
Line-of-sight integration approach to cosmic microwave background anisotropies
ApJ. 469, 437

Hu W., website:
An Introduction to the Cosmic Microwave Background http://background.uchicago.edu/~whu/intermediate/intermediate.html
A Tour of CMB Physics http://background.uchicago.edu/~whu/physics/tour.html
Cosmological Parameters in the CMB http://background.uchicago.edu/~whu/metaanim.html
CMBFAST website:

http://cmbfast.org/

16. Cosmic Microwave Background Polarization

- In addition to CMB temperature anisotropies, polarization of the Microwave Background Signal contains a wealth of information on processes in the early Universe (inflationary gravitational waves and primordial velocity field), on the cosmic matter distribution (weak gravitational lensing of the CMB signal) on secondary postrecombination scattering processes. Of the latter in particular that induced by gas ionized by the first generation of stars and galaxies at and around the epoch of reionization provides an exciting new avenue of research, in particular since the polarization detection by DASI and WMAP. Investigate the physical nature of the various sources of CMB polarization, describe their (expected) imprint (E and B modes) and signature (EE autocorrelation, TE cross-correlation), and the way they can and are exploited to infer important cosmological information.
- Literature suggestions:
 - Bond J.R., 1996
 - Theory and Observations of the Cosmic Background Radiation
 - in "Cosmology and Large Scale Structure", Les Houches LX, eds. R. Schaeffer, J. Silk, M. Spiro, J. Zinn-Justin, Elsevier
 - Challinor A., 2005

Cosmic microwave background polarization analysis

lecture notes, "Data analysis in cosmology", Valencia, 2004

- Hu W., 1995
 - Wandering in the Background: A Cosmic Microwave Background Explorer Ph.D. thesis, Univ. California at Berkeley

website: http://background.uchicago.edu/ \sim whu/thesis/thesispage.html

- Hu W., Dodelson S., 2002
 Cosmic Microwave Background Anisotropies
 ARAA, 40, 171-216
- Hu W., White M., 1997
 A CMB polarization primer
 New Astronomy, 2, 323
- Kamionkowski M., Kosowsky A., Stebbins A., 1997 Statistics of Cosmic Microwave Background Polarization Phys Rev D55, 7368
- Kogut et al., 2003
 Wilkinson Microwave Anisotropy Probe (WMAP) First Year Observations: Temperature Polarization Correlation
 ApJS, 148, 161

Page L., et al., 2003
 First-year WMAP Observations: Interpretation of the TT and TE angular power spectrum peaks
 ApJS, 148, 233

- Page L., et al., 2006
 Three Year Wilkinson Anisotropy Probe (WMAP) Observations: Polarization Analysis
 ApJ, in press
- Seljak U., 1997
 Measuring Polarization in the Cosmic Microwave Background Astrophys. J., 482, 6
- Zaldarriaga M., 1997
 Polarization of the microwave background in reionized models
 Phys. Rev. D., 55, 1822
- Zaldarriaga M., 1998
 Cosmic Microwave Background Polarization Experiments Astrophys. J. 503, 1

- Hu W., website:

- A Polarization Primer
- http://background.uchicago.edu/~whu/polar/webversion/polar.html
- WMAP website: http://map.gsfc.nasa.gov/m_mm/pub_papers/firstyear.html

17. The First Stars

- One of the most important transitions in the state of the Universe is the epoch at which the first stars and galaxies arise out of the gradually unfolding matter distribution. The first generation of stars, Population III stars, are extremely metal poor and will have differed considerably from the stars we know of nowadays. Write an review on the formation and evolution of these stars and investigate the enrichment products following their explosion as supernova.
- Literature suggestions:
 - Abel T., Bryan G.L., Norman M.L., 2002
 The Formation of the First Star in the Universe Science 295, 93
 - Abel T., Bryan G.L., Norman M.L., 2000
 The Formation and Fragmentation of Primordial Molecular Clouds ApJ, 540, 39
 - Alvarez M.A., Bromm V., Shapiro P.R., 2006 The HII region of the first star ApJ 639, 621
 - Barkana R., 2006
 The First Stars in the Universe and Cosmic Reionization
 Science 313, 931
 - Bond J.R., Arnett W.D., Carr B.J., 1984
 The evolution and fate of Very Massive Objects
 ApJ, 280, 825
 - Bromm V., Larson R.B., 2004
 The First Stars
 ARAA 42, 79
 - Bromm V., Coppi P.S., Larson R.B., 2002
 The Formation of the First Stars I. Primordial star-forming cloud
 ApJ, 564, 23
 - Bromm V., Yoshida N., Hernquist L., 2003 The First Supernova Explosions in the Universe ApJ, 596, L135
 - Loeb A., 2006 *First Light* Saas Fee lecture, astro-ph/0603360

 Yoshida N., Omukai K., Hernquist L., Abel T.
 Formation of Primordial Stars in a LCDM Universe ApJ, 652, 6

18. Epoch of Reionization

- One of the most important transitions in the state of the Universe is the epoch at which the first stars and galaxies arise out of the gradually unfolding matter distribution, set ablaze the surrounding gas and very soon thereafter reionizes the hydrogen gas throughout the Universe. The *Epoch of Reionization* is one of the most vigorous and active in today's cosmology, with instruments like LOFAR due to play a seminal role. Discuss in the report the processes and stages in which the reionization of the hydrogen gas proceeds, illustrate this with available computer simulations and assess the shortcomings of the present calculations along with the prospects for detection and possibly mapping of the signal.
- Literature suggestions:
 - Barkana R., 2006
 The First Stars in the Universe and Cosmic Reionization
 Science 313, 931
 - Barkana R., Loeb A., 2006
 The Physics and Early History of the Intergalactic Medium Re. Prog. Phys., 36 pp.
 - Ciardi B., Madau P., 2003
 Probing beyond the epoch of hydrogen reionization with 21 centimeter radiation
 ApJ
 - Ciardi B., Ferrara A., 2004
 The First Cosmic Structures and their Effects
 review, astro-ph/0409018
 - Furlanetto S.R., Sokasian A., Hernquist L., 2003
 Observing the Reionization Epoch Through 21 Centimeter Radiation MNRAS
 - Furlanetto S.R., Peng Oh S., Briggs F.H., 2006 Cosmology at Low Frequencies: the 21cm Transition and the High-Redshift Universe Phys. Rep., 207 pp. On the density of neutral hydrogen in intergalactic space ApJ, 142, 1633
 Gnedin N.Y., Ostriker J.P., 1997
 - Reionization of the Universe and the Early Production of Metals ApJ, 486, 581

- Gnedin N.Y., 2000 Cosmological Reionization by Stellar Sources ApJ 535, 530 - Gnedin N.Y., Fan X., 2006 Cosmic Reionization Redux ApJ, 648, 1 - Illiev I.T. et al., 2006 Reionization observables after WMAP 3-year results MNRAS, astro-ph/0702099 - Loeb A., 2006 First Light Saas Fee lecture, astro-ph/0603360 - Madau P., Meiksin A., Rees M.J., 1997 21 centimeter tomography of the Intergalactic Medium at high redshift ApJ, 475, 429 - Loeb A., Barkana R., 2001 The Reionization of the Universe by the First Stars and Quasars ARAA 39, 19 21 centimeter tomography of the Intergalactic Medium at high redshift ApJ, 475, 429 - Miralda-Escudé J., 2003 The Dark Age of the Universe Science, 300, 1904

19. Inflationary Universe and the Origin of Fluctuations

- The primordial fluctuations out of which structure in the Universe has arisen originated in the inflationary epoch shortly after the Big Bang. Give a review on its basic principles, on the mechanism to generate the fluctuations and the tests of the inflationary scenario via the measurement of the temperature and polarization fluctuations in the cosmic microwave background.
- Literature suggestions:
 - Garcia Y.R., 2005
 - The Origin of the Large-Scale Structure in the Universe: Theoretical and Statistical Aspects
 - Ph.D. thesis, Univ. Lancaster, astro-ph/0507701
 - Guth A., Steinhardt P.J., 1984 *The Inflationary Universe* Scientific American 250, May 1984
 - Kolb E.W., Turner M.S., 1990 *The Early Universe* Frontiers in Physics, Addison-Wesley (you may inform with RvdW).
 - Liddle A.R., 1999 An introduction to cosmological inflation astro-ph/9901124
 - Lineweaver C.H., 2003
 Inflation and the Cosmic Microwave Background
 World Scientific, astro-ph/0305179
 - Page L., et al., 2006
 - Three Year Wilkinson Anisotropy Probe (WMAP) Observations: Polarization Analysis
 - ApJ, in press
 - Peiris H.V., et al., 2003
 First-year WMAP Observations: Implications for Inflation
 ApJS, 148, 233
 - Spergel D.N., et al., 2006 Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology astro-ph

- Tegmark M., 2004 What does inflation really predict astro-ph/0410281