Cosmic Flows

Lecture course University Groningen Feb. 2007-April 2007

Gravitational Instability

$$\frac{\partial \delta}{\partial t} + \frac{1}{a} \nabla \cdot (1 + \delta) \mathbf{v} = 0$$

$$\frac{\partial \mathbf{v}}{\partial t} + \frac{\dot{a}}{a} \mathbf{v} + \frac{1}{a} (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{1}{a} \nabla \phi$$

$$\nabla^2 \phi = 4\pi G \,\bar{\rho} \, a^2 \, \delta(\mathbf{x}, t)$$

Gravitational Instability

$$\frac{\partial \delta}{\partial t} + \frac{1}{a} \nabla \cdot \mathbf{v} = 0$$


$$\frac{\partial \mathbf{v}}{\partial t} + \frac{\dot{a}}{a} \mathbf{v} = -\frac{1}{a} \nabla \phi$$

$$\nabla^2 \phi = \frac{3}{2} \Omega H^2 a^2 \delta(\mathbf{x}, t)$$


Gravitational Instability

$$\frac{\partial^2 \delta}{\partial t^2} \, + \, 2 \frac{\dot{a}}{a} \frac{\partial \delta}{\partial t} \, = \, \frac{3}{2} \Omega_0 H_0^2 \, \frac{1}{a^3} \, \delta$$


Gravitational Instability
$$\delta(\mathbf{x},t) = D_1(t) \Delta_1(\mathbf{x}) + D_2(t) \Delta_2(\mathbf{x})$$




Linear Density Growth
$$D(t) \approx H(t) \int \frac{dt}{a^2 H^2(t)}$$





Gravitational Instability
$$\mathbf{g}(\mathbf{r},t) = -\frac{1}{a}\nabla\phi = \frac{3\Omega H^2}{8\pi}\int\mathrm{d}\mathbf{x}'\delta(\mathbf{x}',t)\frac{(\mathbf{x}'-\mathbf{x})}{|\mathbf{x}'-\mathbf{x}|^3}$$





Local Universe Peculiar Velocities
















Great Attractor














Fig. 16. Distribution in Galactic coordinates of the 76 by Ebeling et al. [129] so far spectroscopically confirmed X-ray clusters (solid dots) of which 80% were previously unknown. Superimposed are Galactic HI column densities in units of 10^{20} cm⁻² (Dickey & Lockman 1990). Note that the region of relatively high absorption $(N_{\rm HI} > 5 \times 10^{21}$ cm⁻²) actually is very narrow and that clusters could be identified to very low latitudes







Fig. 16. Distribution in Galactic coordinates of the 76 by Ebeling et al. [129] so far spectroscopically confirmed X-ray clusters (solid dots) of which 80% were previously unknown. Superimposed are Galactic HI column densities in units of 10^{20} cm⁻² (Dickey & Lockman 1990). Note that the region of relatively high absorption $(N_{\rm HI} > 5 \times 10^{21}$ cm⁻²) actually is very narrow and that clusters could be identified to very low latitudes

PSCz Survey

Dynamics:










PSCz Flow Field


- Flow throughout the volume restored (assuming no shell-crossing)
- ➤ Restored velocities volume-weighted
- ➤ Shot-noise suppressed




PSCz, velocity+density field









Cosmic Dipoles








2MASS survey

- 2MASS all-sky survey: ground-based near-infrared survey whole sky, J(1.2 μ m), H(1.6 μ m), K(2.2 μ m)
- 2MASS extended source catalog (XSC): 1.5 million galaxies
- unbiased sample nearby galaxies
- photometric redshifts: depth in 2MASS maps, "cosmic web" of (nearby) superclusters spanning the entire sky.


courtesy:


T. Jarrett


The Cosmic Web





Looking around us we already see the unmistakable signatures of an intriguing foamlike matter distribution in our immediate Cosmic Vicinity.











Redshift Space Distortions

